


Contents lists available at ScienceDirect

International Journal of Infectious Diseases

journal homepage: www.elsevier.com/locate/ijid

Host biomarkers are associated with progression to dengue haemorrhagic fever: a nested case-control study



Andrea L. Conroy^a, Margarita Gélvez^b, Michael Hawkes^c, Nimerta Rajwans^a,
Vanessa Tran^a, W. Conrad Liles^d, Luis Angel Villar-Centeno^b, Kevin C. Kain^{a,e,*}

^a Sandra A. Rotman Laboratories, Sandra Rotman Centre, University Health Network-Toronto General Hospital, University of Toronto, Toronto, ON, M5G 1L7, Canada

^b Centro de Investigaciones Epidemiológicas, Facultad de Salud, Universidad Industrial de Santander, Bucaramanga, Colombia

^c Department of Pediatrics, University of Alberta, Edmonton, AB, T6G 2E1, Canada

^d University of Washington, Department of Medicine, Seattle, WA, 98195, USA

^e Tropical Disease Unit, Division of Infectious Diseases, Department of Medicine, University of Toronto, Toronto, ON, M5G 2C4, Canada

ARTICLE INFO

Article history:

Received 19 April 2015

Received in revised form 28 July 2015

Accepted 30 July 2015

Corresponding Editor: Eskild Petersen,
Aarhus, Denmark

Keywords:

dengue
biomarkers
pathogenesis
sICAM-1
soluble endoglin
CXCL10

ABSTRACT

Objectives: Dengue represents the most important arboviral infection worldwide. Onset of circulatory collapse can be unpredictable. Biomarkers that can identify individuals at risk of plasma leakage may facilitate better triage and clinical management.

Design: Using a nested case-control design, we randomly selected subjects from a prospective cohort study of dengue in Colombia ($n = 1582$). Using serum collected within 96 hours of fever onset, we tested 19 biomarkers by ELISA in cases (developed dengue hemorrhagic fever or dengue shock syndrome (DHF/DSS); $n = 46$), and controls (uncomplicated dengue fever (DF); $n = 65$) and healthy controls (HC); $n = 15$). **Results:** Ang-1 levels were lower and angptl3, sKDR, sEng, sICAM-1, CRP, CXCL10/IP-10, IL-18 binding protein, CHI3L1, C5a and Factor D levels were increased in dengue compared to HC. sICAM-1, sEng and CXCL10/IP-10 were further elevated in subjects who subsequently developed DHF/DSS ($p = 0.008$, $p = 0.028$ and $p = 0.025$, respectively). In a logistic regression model, age (odds ratio (OR) (95% CI): 0.95 (0.92–0.98), $p = 0.001$), hyperesthesia/hyperalgesia (OR; 3.8 (1.4–10.4), $p = 0.008$) and elevated sICAM-1 ($>298\text{ng/mL}$: OR; 6.3 (1.5–25.7), $p = 0.011$) at presentation were independently associated with progression to DHF/DSS.

Conclusions: These results suggest that inflammation and endothelial activation are important pathways in the pathogenesis of dengue and sICAM-1 levels may identify individuals at risk of plasma leakage. © 2015 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Background

Dengue is an emerging infectious disease of global importance with 2.5 to almost 4 billion people residing in dengue endemic countries in which an estimated 50–100 million infections occur annually.^{1,2} Historically, the highest risk areas for dengue fever were in Southeast Asia and the western Pacific (where

approximately 75% of the global disease burden is located). However, the risk of dengue, and severe dengue, is increasing in the Americas following disease re-emergence after lapsed vector control programs.³ Between 2000 and 2006, the majority of cases reported (68%) were registered in this region.⁴ Cyclical outbreaks of dengue in the Caribbean, and Central and South America have resulted in considerable morbidity and mortality. In 2013, 2.35 million cases of dengue were reported in the Americas, of which 37,687 cases were classified as severe dengue, with Brazil and Colombia recording the highest incidence.

Dengue is an important consideration in the differential diagnosis of fever in endemic countries and in travelers returning from these regions.⁵ Dengue infection presents with a wide spectrum of signs and symptoms, including rapid onset of fever that may be accompanied by myalgia, arthralgia, retro-orbital pain, facial flushing, erythema, nausea, vomiting, cutaneous

* Corresponding author. Sandra Rotman Laboratories, Sandra Rotman Centre for Global Health, MaRS Centre, Toronto Medical Discovery Tower, Suite 10-360A, 101 College St., Toronto, ON, M5G 1L7, Canada. Tel.: +14165817704.

E-mail addresses: andrea.conroy@utoronto.ca (A.L. Conroy), margarita.gelvez@hotmail.com (M. Gélvez), mthawkes@ualberta.ca (M. Hawkes), nrajwans@uhnresearch.ca (N. Rajwans), vanessa.tran@utoronto.ca (V. Tran), WCLiles@medicine.washington.edu (W.C. Liles), luisangelvillarc@gmail.com (L.A. Villar-Centeno), kevin.kain@uhn.ca (K.C. Kain).

hyperesthesia or hyperalgesia, and minor bleeding abnormalities as evidenced by a positive tourniquet test. A small percentage of individuals with dengue will develop life-threatening complications, which can include plasma leak and circulatory collapse. The clinical evolution of disease can be unpredictable making the initial clinical assessment a critical step in patient management.⁶

Dengue is a single-stranded RNA virus with four distinct serotypes (DEN-1 to -4). Risk of haemorrhage and plasma leak is generally higher in secondary infections, but other factors including age, sex, and other host and viral determinants also contribute to disease susceptibility. The prevailing hypothesis is that cross-reactive, non-neutralizing antibodies in secondary heterotypic infections facilitate viral entry into target cells, thereby increasing viral load, enhancing deleterious inflammatory responses and contributing to endothelial activation and dysfunction (reviewed in⁷). The endothelium is central to dengue pathogenesis as increased vascular permeability and fragility are defining features in severe disease. However, inflammation, coagulation and complement activation are also involved in disease pathogenesis, and alterations in soluble mediators from these pathways have been described in severe dengue infections.^{8–18}

Clinical management of dengue depends on appropriate triage, referral and treatment of cases, especially in outbreaks. Although there are no specific anti-infective or immunomodulatory therapies to prevent plasma leak, appropriate fluid management and supportive care can reduce the mortality rate from over 20% to less than 1%.^{3,19} With rates of severe disease on the rise, especially in the Americas,^{20–22} new tools are urgently needed to improve patient triage and risk prediction of severe disease.²³ We investigated whether host biomarkers, chosen from pathways implicated in dengue pathogenesis, were altered in dengue relative to healthy controls, and whether the biomarkers assessed during the acute phase of disease were associated with the subsequent development of dengue haemorrhagic fever (DHF) or dengue shock syndrome (DSS).

2. Methods

2.1. Study Population

All four dengue virus serotypes circulate in Colombia, which reports the highest number of severe dengue cases and deaths in the region.^{3,24} Bucaramanga is a metropolitan area in northeastern Colombia where the incidence of dengue ranges from 113–269 cases per 100,000 people.^{25–27} Participants from Bucaramanga who presented within 96 hours of fever onset were eligible for recruitment if they were greater than five years of age. Subjects were excluded based on the presence of the following conditions: history of concomitant diseases such as diabetes, acquired immunodeficiency syndrome (AIDS), hematologic disorders, cancer, or cardiac disease, albuminemia (<3 g/dL), evidence of severe dengue at presentation, for example major bleeding, effusions, or shock. After participants provided informed consent, a physical examination was performed, and a blood sample was obtained to determine hematocrit, albumin levels, and platelet and leukocyte counts. A serum sample was collected and stored at -80°C for future biomarker assessment.

Diagnosis of dengue virus infection was confirmed based on viral isolation or serology (seroconversion from a negative to a positive IgM test or a four-fold increase in dengue antibodies in a convalescent blood sample). All study participants were enrolled before the development of DHF/DSS. Participants were followed daily until day 7 of disease with daily microhematocrit measurements to facilitate the recognition of severe dengue. Platelet counts were repeated daily for subjects with previous platelet counts less

than 120,000/mm³ or if there were signs of spontaneous haemorrhage, effusion, oedema or change in hematocrit > 10%. Study participants were classified as DF or DHF/DSS following recovery and full chart review using the 1997 WHO classification system according to the study design.²⁸

2.2. Study Design

This study was a case-control study nested within the prospective cohort study (n = 1582) of suspected dengue in Bucaramanga, Colombia. Blood samples were collected from study participants at clinical presentation. Cases were individuals who developed DHF/DSS and controls had dengue fever (DF). Study subjects with available serum samples were eligible for biomarker testing and were randomly chosen from the cohort database using computer generated simple randomization. Serum samples were collected from 15 healthy Bucaramangan adults to derive a population-based normal range.

2.3. Biomarker Assessment

Serum concentrations of biomarkers were measured in samples collected at presentation using ELISA DuoSets from R&D Systems (Minneapolis, MN). Biomarkers measured were C-reactive protein (CRP), CXCL10/IP-10, IL-18 binding protein (IL-18BP), IL-10, chitinase 3-like 1 (CHI3L1), C5a, complement factor D, angiopoietin-1 (Ang-1), angiopoietin-2 (Ang-2), soluble Tie-1 (sTie-1), soluble Tie-2 (sTie-2), angiopoietin-like protein 3 (Angptl3), angiopoietin-like protein 4 (Angptl4), vascular endothelial growth factor (VEGF), soluble VEGF R1 (sFlt-1), soluble VEGF R2 (sKDR), soluble endoglin (sEng), soluble intercellular adhesion molecule-1 (sICAM-1) and platelet factor 4. All ELISAs were validated prior to use, and sample dilutions were optimized for each biomarker using a dilution curve of serum obtained from febrile subjects with dengue fever in Bucaramanga. ELISAs were tested according to the manufacturer's instructions (R&D Systems) with minor modifications previously described.²⁹

2.4. Statistical Analysis

GraphPad Prism v5, SPSS v20 and MedCalc® v12 were used for statistical analysis. Comparisons of continuous variables were performed using the Mann-Whitney U test. Comparisons of proportions were performed using Pearson chi-square test or Fisher's exact test, as appropriate. Exploratory logistic regression models were built using forward step-wise selection including all variables with $p < 0.10$ by bivariate analysis. Model fit was assessed using the Hosmer-Lemeshow test and ensuring the term was not significant $p > 0.05$. Biomarkers were dichotomized by generating a ROC curve and using the Youden index to identify the optimal cut-off (Youden index: $J = \max [\text{sensitivity} + \text{specificity}] - 1$).

3. Results

3.1. Clinical Characteristics of Subjects that Developed DHF/DSS

111 subjects with microbiologically confirmed dengue virus infection were included in the current study along with 15 healthy controls from Bucaramanga. 65 had DF while 46 developed DHF/DSS over the course of clinical follow up.³ Individuals that developed DHF/DSS were significantly younger, had higher axillary temperatures and lower mean arterial pressure at presentation and were more likely to report chills, hyperesthesia/hyperalgesia, and dizziness than those with DF (Table 1). The median day of illness at presentation was day 4 for both groups ($p = 0.461$,

Table 1
Demographic and clinical features of cohort at presentation

	DF (n = 65)	DHF/DSS (n = 46)	P value
Demographics			
Age, years	28.0 (19.0–45.5)	21.0 (14.0–31.3)	0.006
Sex (% F)	35 (53.8)	25 (54.3)	0.958
Height, cm	163 (155–172)	160 (150–168)	0.094
Weight, kg	63.0 (54.0–76.5)	59.0 (47.0–70.0)	0.156
Duration of fever, hours	77.0 (66.8–89.0)	75.3 (60.1–88.8)	0.733
Laboratory			
Axillary temperature, °C	36.3 (36.0–37.0)	36.8 (35.9–37.9)	0.067
Platelet count ($\times 10^3 \mu\text{L/mL}$)	126.0 (83.0–182.0)	119.5 (79.8–176.3)	0.940
Leukocyte count	2900 (2100–4000)	2950 (2275–3925)	0.682
Hematocrit	39.4 (36.1–42.6)	38.0 (35.5–42.0)	0.436
Mean arterial pressure	88.0 (80.7–94.0)	83.7 (76.7–90.8)	0.038
Tourniquet test (# petechiae)	22.0 (4.5–32.5)	17.5 (5.0–25.0)	0.281
Aspartate aminotransferase ^a	59.9 (40.0–114.8)	56.5 (43.0–95.0)	0.958
Alanine aminotransferase ^a	37.6 (22.0–57.0)	30.7 (24.0–74.0)	0.889
Lactate dehydrogenase ^b	320.0 (200.5–433.1)	333.8 (172.0–716.9)	0.540
Albumin ^c	4.5 (4.2–4.7)	4.1 (3.8–4.5)	0.069
Creatinine phosphokinase ^c	134.6 (59.8–769.0)	76.6 (54.1–184.5)	0.170
Cholesterol ^d	155.1 (141.0–164.0)	148.6 (113.3–154.6)	0.118
Triglycerides ^d	121.7 (87.2–175.2)	159.1 (72.7–203.3)	1.000
Signs and Symptoms (%)			
Headache	61 (93.8)	44 (95.7)	0.678
Retro-orbital pain	43 (66.2)	35 (76.1)	0.259
Asthenia	46 (70.8)	31 (67.4)	0.704
Muscle Pain	59 (90.8)	41 (91.1), n = 45	0.951
Joint Pain	52 (80.0)	36 (80.0), n = 45	1.000
Chills	64 (98.5)	41 (89.1)	0.032
Sore throat	23 (35.4)	12 (28.3)	0.430
Facial erythema	35 (53.8)	22 (47.8)	0.532
Rash	25 (38.5)	23 (53.5), n = 43	0.124
Hyperesthesia/hyperalgesia	15 (23.1)	17 (40.5)	0.055
Nausea	47 (72.3)	38 (82.6)	0.207
Vomiting	20 (30.8)	18 (39.1)	0.360
Diarrhea	25 (38.5)	16 (34.8)	0.692
Abdominal pain	31 (48.4)	30 (65.2)	0.081
Blurred vision	24 (36.9)	20 (44.4)	0.429
Dizziness	39 (60.0)	35 (77.8)	0.051
Conjunctival injection	31 (47.7)	17 (37.0)	0.261
Orthostatic hypotension	8 (12.3)	9 (19.6)	0.296

Participants with dengue fever (DF) that did not develop complications, DF participants that developed DHF/DSS.

^a DF (n = 40), DHF/DSS (n = 31); ^b DF (n = 38), DHF/DSS (n = 31); ^c DF (n = 24), DHF/DSS (n = 18); ^d DF (n = 15), SD (n = 13).

chi-square) and the median time to DHF/DSS diagnosis was 2 days after presentation.

3.2. Alterations of Host Biomarkers in Dengue

We evaluated 19 host biomarkers in patients with dengue virus infection and compared them to healthy controls to determine which markers were elevated during the acute febrile phase of dengue (2–4 days after symptom onset). Biomarkers were broadly chosen from four pathways implicated in dengue pathogenesis: inflammation, endothelial activation, complement activation, and coagulation. Eleven biomarkers differed in dengue infection compared to healthy controls ($p < 0.05$ by Mann-Whitney U test, Table 2). Ang-1 was significantly reduced while Angptl3, sKDR, sEng, sICAM-1, CRP, CXCL10 (IP-10), IL-18 binding protein, CHI3L1, C5a and Factor D were increased in dengue fever compared to healthy controls (Figure 1).

3.3. Host Biomarkers are Associated with the Development of Hemorrhage

We hypothesized that host biomarkers altered in dengue precede clinical manifestations of hemorrhage and vascular leak. Three biomarkers were significantly elevated at presentation in subjects that subsequently developed DHF/DSS compared to those with an uncomplicated disease course: sICAM-1, sEng and CXCL10 ($p = 0.008$, $p = 0.028$ and $p = 0.025$ respectively) (Table 2, Figure 2).

Levels of CXCL10 were significantly correlated with sICAM-1 levels (Spearman's rho, p -value; 0.468, $p < 0.001$) and sEng levels (0.348, $p < 0.0001$), but there was no statistically significant association between sEng and sICAM-1 (0.184, $p = 0.053$).

We explored the relationship between the biomarkers associated with disease progression and routine clinical laboratory values at the time of clinical presentation, including platelet and leukocyte counts, hematocrit, bleeding tendency, mean arterial pressure and liver enzymes. Levels of sICAM-1 and sEng correlated weakly and negatively with the platelet count (Spearman's rho, p -value: sICAM-1, -0.197 , $p = 0.038$; sEng, -0.219 , $p = 0.021$). CXCL10 correlated with both platelet and leukocyte counts (platelet count, -0.533 , $p < 0.0001$; leukocyte count, -0.355 , $p < 0.0001$). Biomarker levels were not correlated with hematocrit, the number of petechiae by tourniquet test or mean arterial pressure. Both sICAM-1 and CXCL10 were significantly correlated with AST and ALT (AST: sICAM-1, 0.375, $p = 0.001$; CXCL10, 0.314, $p = 0.008$; ALT: sICAM-1, 0.410, $p < 0.0001$; CXCL10, 0.287, $p = 0.015$). sEng did not correlate with AST or ALT but was negatively correlated with total cholesterol and low density lipoproteins (cholesterol, -0.465 , $p = 0.013$; low density lipoprotein, -0.601 , $p = 0.001$).

3.4. Predictive Models of DHF: Integrating Clinical and Biomarker Data

Based on the hypothesis that host biomarkers can add value to clinical findings, we explored models to integrate our biomarker

Table 2
Biomarkers in Dengue

	HC (n = 15)	Dengue	
		DF (n = 65)	DHF/DSS (n = 46)
Inflammation			
CRP ^a	1.8 (2.3–6.1)***	18.7 (9.2–41.3)	21.0 (6.4–41.4)
IP-10	0.2 (0.2-0.4) ***	2.8 (1.0–4.4)	3.6 (1.8–5.0) †
IL-18 BP	5.8 (4.3–8.3) ***	62.3 (36.4–97.0)	75.9 (52.8–90.6)
IL-10 ^b	0.03 (0.02-0.2)	145.2 (19.5–390.9)	58.0 (19.5–357.9)
CHI3L1	44.5 (35.8–61.3)*	61.6 (44.6–81.2)	57.3 (40.3–78.3)
Complement System			
C5a	18.5 (13.8–22.1) ***	53.8 (33.9–62.8)	52.5 (35.3–67.4)
Factor D ^a	1.1 (1.0–1.3)**	1.4 (1.1–1.6)	1.3 (1.2–1.6)
Endothelium			
Ang-1	44.9 (33.0–53.6)*	32.2 (25.6–43.9)	31.6 (22.3–47.0)
Ang-2	1.4 (1.0–2.6)	1.7 (1.3–2.4)	2.1 (1.3–2.9)
sTie-1	8.3 (6.8–12.7)	10.5 (8.1–12.9)	9.6 (8.1–13.5)
sTie-2	8.7 (6.4–9.7)	7.3 (5.5–8.9)	8.1 (6.0–9.9)
Angptl3	108.4 (82.3–117.5) ***	164.0 (140.6–192.9)	170.8 (146.5–196.3)
Angptl4	44.7 (30.2–90.9)	37.8 (29.3–56.3)	41.7 (33.6–64.2)
VEGF ^b	0.2 (0.2-0.3)	128.2 (48.8–291.9)	124.3 (48.8–321.5)
sFlt-1 ^b	39.1 (39.1–343.5)	39.1 (39.1–863.3)	94.6 (39.1–1074)
sKDR	5.3 (4.5–6.0) *	6.1 (5.5–7.1)	6.1 (5.5–7.1)
sEng	6.6 (5.7–8.4) ***	11.0 (9.2–13.2)	12.8 (10.3–14.7) †
sICAM-1	169.3 (135.8–287.3) ***	329.1 (284.1–467.1)	399.9 (323.0–539.0) ††
Coagulation			
Platelet factor 4 ^a	18.5 (15.6–27.1)	25.7 (16.2–41.2)	23.8 (15.6–37.7)

Healthy controls (HC), Participants with dengue that did not develop complications (DF), participants with dengue that developed DHF/DSS (DHF/DSS).

Biomarker concentration in ng/mL unless otherwise indicated, ^a µg/mL, ^b pg/mL.

*** $p < 0.001$ (HC vs. all dengue), ** $p < 0.01$ (HC vs. all dengue), * $p < 0.05$ (HC vs. all dengue).

†† $p < 0.01$ (DF vs. DHF/DSS), † $p < 0.05$ (DF vs. DHF/DSS).

findings with clinical data. All variables with $p < 0.10$ from bivariate analysis (Table 1) were considered for the predictive models and biomarkers were included as dichotomous variables with the following cut-offs (CXCL10 > 1.07 ng/mL; sEng > 11.8 ng/mL; sICAM-1 > 298.2 ng/mL).

Using forward step-wise logistic regression, we developed a clinical model consisting of age and hyperesthesia/hyperalgesia (Table 3). We then added the biomarkers into the clinical model to generate a biomarker model that included age, hyperesthesia/hyperalgesia and sICAM-1. In the biomarker model, age was associated with a 0.95 (95% CI, 0.92–0.98, $p = 0.002$) odds of DHF/DSS while the presence of hyperesthesia/hyperalgesia was associated with 3.8-fold increased odds of developing DHF/DSS (95% CI, 1.4–10.4, $p = 0.008$). sICAM-1 > 298.2 ng/mL was associated with a 6.3-fold increased odds of developing DHF/DSS (95% CI, 1.5–25.7, $p = 0.011$) (Table 3).

4. Discussion

Dengue fever is an emerging infectious disease endemic to the tropics and sub-tropics that causes cyclical outbreaks or epidemics that can overwhelm health care resources. Only a small proportion of infections will progress to severe disease (DHF/DSS), but it is difficult to accurately identify individuals at risk of plasma leak and circulatory collapse during the acute phase of disease when individuals are likely to first seek medical care. In this study, we investigated whether host proteins derived from pathways implicated in the pathogenesis of dengue would differentiate between subjects with an uncomplicated disease course versus those that developed DHF/DSS. Compared to healthy controls, eleven proteins were altered in dengue infection (ang-1, angptl3, sKDR, sEng, sICAM-1, CRP, CXCL10/IP-10, IL-18BP, CHI3L1, C5a, Factor D), and three proteins were significantly elevated at presentation in individuals who subsequently developed DHF/DSS (sEng, sICAM-1 and CXCL10). sICAM-1 was an independent predictor of DHF/DSS (Figure 2, Table 3). These results suggest that host biomarkers, once validated, may have clinical utility in patient

triage and risk-stratification for referral, admission and implementation of timely supportive measures.

Consistent with other studies, levels of sICAM-1, CRP, CXCL10, C5a and Factor D were elevated and levels of Ang-1 were reduced in subjects with dengue.^{11,12,16,30–36} IL-10 was neither elevated in dengue nor was it associated with the development of DHF/DSS in this population unlike findings reported from Venezuela,^{37,38} but consistent with a recent study from Sri Lanka.³⁹ There were no differences in levels of VEGF or its receptors in predicting DHF/DSS, unlike a report from Thailand.⁴⁰ These discrepancies may reflect age differences;⁴¹ however, other factors (host and viral genotype or the kinetics of infection) may also be involved.

This study identified four novel biomarkers associated with dengue infection: CHI3L1, IL-18 BP, Angptl3, and sEng. CHI3L1 is expressed by a variety of cells and tissues (including leukocytes and endothelium) in response to inflammatory stimuli and is involved in modulating inflammatory responses (reviewed in⁴²). Elevated levels of CHI3L1 have been associated with dysregulated inflammatory responses in other infections such as pneumonia and malaria.^{43–45} IL-18BP binds and neutralizes IL-18, a pro-inflammatory cytokine that is induced by and further induces expression of IFN- γ . IL-18 has been implicated in resistance to dengue infection in a murine model,⁴⁶ but IL-18 has also been associated with dengue disease severity in humans.¹⁷ Angptl3 is a protein expressed by the liver and is involved in regulating angiogenesis and lipid metabolism;^{47–49} however, Angptl3 levels were not associated with levels of cholesterol (total, HDL or LDL) triglycerides or liver enzymes AST or ALT in this study.

Endoglin (Eng) is an integral membrane protein that serves as a co-receptor for protein members of the transforming growth factor (TGF)- β family. It is predominantly expressed on endothelium and is highly expressed at sites of active angiogenesis (including inflammation and vascular injury).^{50,51} sEng is generated through proteolytic processing of membrane bound Eng through matrix metalloproteinase-14⁵² and regulates vascular tone and permeability in addition to modulating immune responses.⁵³ Overexpression of sFlt-1 and sEng leads to increased microvascular




Figure 1. Host biomarkers that are altered in dengue infection.

(A–K) Box and whisker plots showing the distribution of biomarkers significantly different ($p < 0.05$) in healthy controls from Bucaramanga ($n = 15$) compared to subjects with serological confirmation of dengue fever ($n = 111$). Boxes show the median and interquartile range while the whiskers denote the minimum/maximum greatest value excluding outliers (defined as values 1.5x greater than the 3rd quartile or 1.5x less than the 1st quartile). The mean is denoted by the plus-sign (+). Data were analysed using the Mann-Whitney U test.




Figure 2. Biomarkers associated with the development of DHF/DSS.

Median and scatter of serum biomarkers (sICAM-1, sEng, CXCL10) measured at presentation in dengue infection in participants with uncomplicated dengue fever (DF, n = 65) and those who developed DHF/DSS (n = 46). Data were analysed using Mann-Whitney U test.

permeability in vivo and has an additive effect on increasing microvascular permeability in an organ specific manner.⁵³ Elevated sEng levels have been reported in a number of infectious and non-infectious conditions associated with vascular dysfunction including malaria and preeclampsia.^{53–57} In this study, sEng levels were higher in individuals that subsequently developed DHF/DSS compared to those with DF. These findings are consistent with reports of sEng as a vascular permeability factor, but conflict with a recent study that found no difference in circulating sEng levels in dengue patients with plasma leakage assessed by X-ray or ultrasound on admission.⁵⁸ It is also possible that sEng may be modulating inflammatory responses through its effects on TGF- β signalling.

CXCL10 is an IFN- γ induced proinflammatory chemokine that is highly expressed by primary cells infected with dengue virus^{34,59,60} and by primary cells from dengue-infected patients.⁶¹ CXCL10 promotes viral clearance by binding to heparin sulfate, a co-receptor for viral entry,^{62,63} and peak levels of CXCL10 have been temporally associated with maximal virus production from mononuclear cells.⁶⁴ In murine models, CXCL10 is associated with resistance to dengue infection.^{63,65} CXCL10 levels are elevated during the acute febrile phase of infection^{34,35,66} and are higher in dengue than other febrile illnesses.^{33,34} There are little human data on CXCL10 levels and disease severity. A report from Colombia noted increased expression of CXCL10 in the cardiac tissue of a fatal case of dengue haemorrhagic fever,⁶⁷ but this finding requires validation. In this study, CXCL10 levels were higher in individuals that developed DHF/DSS. It is possible that CXCL10 was upregulated as an anti-viral response and individuals that developed DHF/DSS had higher viral loads, consistent with other reports.^{68–70} Further studies are required to determine whether CXCL10 in humans is associated with resistance by reducing viral load or whether it may exacerbate disease by amplifying proinflammatory responses.

ICAM-1, a member of the immunoglobulin superfamily, exists as both a transmembrane (ICAM-1) and a soluble protein (sICAM-1). ICAM-1 is expressed on a variety of cells including leukocytes, hepatocytes and endothelial cells and is upregulated by inflammatory stimuli. Increased expression of ICAM-1 has been reported in dengue fever on endothelium^{18,71–74} and leukocytes.^{75,76} Upregulation of endothelial ICAM-1 occurs upon stimulation with sera from dengue patients, supernatants from dengue-infected cells or via direct stimulation with virus antigen in vitro. Stimulation occurs, at least in part, in a TNF and NF- κ B-dependent manner.^{18,71–73} Elevated sICAM-1 levels have been reported in dengue fever but have not been previously associated with disease severity,^{30,31,75,77} except in a study by Bethell et al., where sICAM-1 was decreased in Vietnamese children with dengue shock syndrome. In the latter study, the decrease was attributed to a loss of protein secondary to capillary leak that occurs in severe dengue.⁷⁸

In this study, sICAM-1 levels were increased in individuals with dengue infection compared to healthy controls (Figure 1), and sICAM-1 levels were higher in individuals that subsequently developed DHF/DSS compared to individuals with an uncomplicated course of DF (Figure 2). This effect was most pronounced when samples were collected prior to defervescence. sICAM-1 levels were positively correlated with liver enzymes AST and ALT and negatively correlated with platelet counts. Decreased platelet counts and increases in AST have been associated with severe disease onset in dengue fever,^{79,80} but were not predictive in this nested case-control study, which may reflect a lack of power. sICAM-1 was an independent predictor of DHF/DSS in a clinical model alongside age and hyperesthesia/hyperalgesia and improved clinical prediction. These findings suggest that early systemic inflammation in dengue may lead to upregulation of cell-bound ICAM-1 and trigger cleavage of sICAM-1. The kinetics and functional consequences of elevated sICAM-1 in dengue infection require further investigation.

Table 3
Clinical and Biomarker Logistic Regression Models to Predict DHF/DSS

	B (S.E.)	Wald	Df	P Value	OR (95% CI)
CLINICAL					
Age	-0.047 (0.015)	9.397	1	.002	0.95 (0.93–0.98)
Hyperesthesia/hyperalgesia	1.184 (0.480)	6.088	1	.014	3.3 (1.3–8.4)
Constant	0.471 (0.432)	1.185	1	.276	1.6
BIOMARKER					
Age	-.052 (0.016)	10.733	1	.001	0.95 (0.92–0.98)
Hyperesthesia/hyperalgesia	1.345 (0.511)	6.933	1	.008	3.8 (1.4–10.4)
sICAM-1 (>298.2ng/mL)	1.837 (0.719)	6.523	1	.011	6.3 (1.5–25.7)
Constant	-0.998 (0.748)	1.782	1	.182	0.37

Coefficient, B; standard error, (S.E.); degrees of freedom, Df; odds ratio, OR; confidence interval, CI.

CLINICAL: Hosmer-Lemeshow (Chi-square, 12.506; df, 7; p=0.085), -2 Log Likelihood (128.383), Cox & Snell R² (0.131).

BIOMARKER: Hosmer-Lemeshow (Chi-square, 11.008; df, 8; p=0.201), -2 Log Likelihood (120.058), Cox & Snell R² (0.196).

Dengue has significant health, economic and social impacts on affected communities.⁸¹ On average, a hospitalized case of dengue fever costs three times that of an ambulatory case,⁸² and tools that can facilitate better patient triage at time of presentation with fever may reduce unnecessary hospital admissions and support the rational allocation of resources. A number of clinical signs and symptoms, laboratory parameters and biochemical markers have been associated with increased risk of clinical deterioration in dengue infection. However, the predictive accuracy of individual parameters varies according to the population studied and epidemiologic risk factors, and viral genotype. Host genetics may also affect host susceptibility to severe disease. The majority of published papers focus on paediatric populations from Southeast Asia; less is known about disease progression and markers of disease severity in adult populations and populations from Latin America. As Colombia has one of the highest rates of severe dengue and associated mortality in the western hemisphere^{24,83}, it is important to identify markers that can facilitate improved patient triage in this setting. In our study we found that patient age and the presence of cutaneous hyperesthesia/hyperalgesia, an under-recognized clinical symptom, were independently associated with progression to DHF/DSS.

Of interest, two candidate biomarkers associated with DHF/DSS progression in this study (CXCL10 and sICAM-1), are associated with disease severity in other viral haemorrhagic fevers. Increases in serum sICAM-1 and CXCL10 are associated with Crimean-Congo haemorrhagic fever (CCHF), and haemorrhagic fever with renal syndrome,^{84–87} and sICAM-1 has been identified as a predictive marker of CCHF mortality.⁸⁴ Serum CXCL10 has been found to be significantly higher in patients with fatal yellow fever and fatal Zaire virus-associated Ebola virus disease (EVD); however, the prognostic value was not assessed in these studies.^{88,89} Increases in sICAM-1 have been associated with fatal outcomes in Ugandan children with Sudan virus-associated EVD infection.⁹⁰ In the same outbreak, both sICAM-1 and CXCL10 were elevated in patients with haemorrhage.⁹¹ These data suggest that the biomarker candidates identified in our cohort may also identify patients at risk of complications in other viral infections associated with endothelial pathobiology and haemorrhage.

This study represents one of the larger studies to assess the association of biomarkers with disease progression in samples prospectively collected during acute disease. Although the results are encouraging, they require validation in a larger prospective study and longitudinal sampling over the course of illness to evaluate biomarker kinetics. These findings also need to be validated using the new WHO dengue classification system to determine how the biomarkers relate to warning signs, organ dysfunction, and the development of severe dengue. Although further mechanistic studies are required to elucidate the role, if any, of the markers on mediating disease, our findings suggest there may be specific immune signatures that precede clinical deterioration. Further investigations into these pathways may identify novel markers to assist in patient triage, and inform the design of new interventions.

Acknowledgments

The authors wish to express their appreciation to all of the study participants and study staff.

Competing Interests: The authors declare that they have no competing interests.

Authors' Contributions: ALC, MG, MH, NR, VT, WCL, LAVC and KCK participated in the design of the study. ALC, MG and NR carried out the immunoassays. ALC and MH performed the statistical analysis. WCL, LAVC and KCK contributed reagents/ materials/ analysis tools. ALC, MH, VT and KCK wrote the initial draft of the

manuscript. MG, NR, VT, WCL and LAVC critically revised the manuscript. All authors read and approved the final manuscript.

Funding: This work was supported by Colciencias [1102-459-21561 to LAVC] and the Canadian Institutes of Health Research [MOP-115160, MOP-136813 and MOP-13721 to KCK; Canada Research Chairs to KCK and WCL; Clinician Scientist Training Award to MH; Fellowship award to ALC]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Ethics Statement: Ethical approval for this study was granted from the Medical Ethics Committee of the Universidad Industrial de Santander in Bucaramanga, Colombia. All subjects (or parents/guardians for minors) provided written informed consent to enter the study. Selection of subjects for biomarker testing and subsequent biomarker data analysis was anonymous.

References

- WHO. Dengue and severe dengue. In: WHO Media centre; 2014.
- Brady OJ, Gething PW, Bhatt S, et al. Refining the Global Spatial Limits of Dengue Virus Transmission by Evidence-Based Consensus. *PLoS Negl Trop Dis* 2012;**6**:e1760.
- WHO. Dengue Guidelines for Diagnosis, Treatment, Prevention and Control. World Health Organization; 2009.
- San Martin JL, Brathwaite O, Zambrano B, et al. The epidemiology of dengue in the Americas over the last three decades: a worrisome reality. *Am J Trop Med Hyg* 2010;**82**:128–35.
- Schwartz E, Weld LH, Wilder-Smith A, et al. Seasonality, annual trends, and characteristics of dengue among ill returned travelers, 1997–2006. *Emerging infectious diseases* 2008;**14**:1081–8.
- Yacoub S, Wills B. Predicting outcome from dengue. *BMC medicine* 2014;**12**:147.
- Halstead SB. Neutralization and antibody-dependent enhancement of dengue viruses. *Advances in virus research* 2003;**60**:421–67.
- Huang YH, Liu CC, Wang ST, et al. Activation of coagulation and fibrinolysis during dengue virus infection. *J Med Virol* 2001;**63**:247–51.
- Suharti C, van Gorp EC, Setiati TE, et al. The role of cytokines in activation of coagulation and fibrinolysis in dengue shock syndrome. *Thrombosis and haemostasis* 2002;**87**:42–6.
- Sosothikul D, Seksarn P, Pongsewalak S, Thisyakorn U, Lusher J. Activation of endothelial cells, coagulation and fibrinolysis in children with Dengue virus infection. *Thrombosis and haemostasis* 2007;**97**:627–34.
- Nascimento EJ, Silva AM, Cordeiro MT, et al. Alternative complement pathway deregulation is correlated with dengue severity. *PLoS One* 2009;**4**:e6782.
- Avirutnan P, Punyadee N, Noisakran S, et al. Vascular leakage in severe dengue virus infections: a potential role for the nonstructural viral protein NS1 and complement. *J Infect Dis* 2006;**193**:1078–88.
- Priyadarshini D, Gadia RR, Tripathy A, et al. Clinical findings and pro-inflammatory cytokines in dengue patients in Western India: a facility-based study. *PLoS One* 2010;**5**:e8709.
- Houghton-Trivino N, Salgado DM, Rodriguez JA, Bosch I, Castellanos JE. Levels of soluble ST2 in serum associated with severity of dengue due to tumour necrosis factor alpha stimulation. *The Journal of general virology* 2010;**91**:697–706.
- Seet RC, Chow AW, Quek AM, Chan YH, Lim EC. Relationship between circulating vascular endothelial growth factor and its soluble receptors in adults with dengue virus infection: a case-control study. *International journal of infectious diseases: IJID: official publication of the International Society for Infectious Diseases* 2009;**13**:e248–53.
- Malasit P. Complement and dengue haemorrhagic fever/shock syndrome. *The Southeast Asian journal of tropical medicine and public health* 1987;**18**:316–20.
- Mustafa AS, Elbishbishi EA, Agarwal R, Chaturvedi UC. Elevated levels of interleukin-13 and IL-18 in patients with dengue haemorrhagic fever. *FEMS immunology and medical microbiology* 2001;**30**:229–33.
- Anderson R, Wang S, Osioy C, Issekutz AC. Activation of endothelial cells via antibody-enhanced dengue virus infection of peripheral blood monocytes. *J Virol* 1997;**71**:4226–32.
- Hung NT. Fluid management for dengue in children. *Paediatrics and international child health* 2012;**32**(Suppl 1):39–42.
- Guzman MG, Kouri G. Dengue and dengue hemorrhagic fever in the Americas: lessons and challenges. *J Clin Virol* 2003;**27**:1–13.
- Guha-Sapir D, Schimmer B. Dengue fever: new paradigms for a changing epidemiology. *Emerg Themes Epidemiol* 2005;**2**:1.
- Harris E, Vide E, Perez L, et al. Clinical, epidemiologic, and virologic features of dengue in the 1998 epidemic in Nicaragua. *Am J Trop Med Hyg* 2000;**63**:5–11.
- Simmons CP, Farrar JJ, Nguyen V, Wills B. Dengue. *N Engl J Med* 2012;**366**:1423–32.
- Villar LA, Rojas DP, Besada-Lombana S, Sarti E. Epidemiological trends of dengue disease in Colombia (2000–2011): a systematic review. *PLoS Negl Trop Dis* 2015;**9**:e0003499.
- SIVIGILA. Comportamiento por regiones del dengue en el 2001. Boletín Epidemiológico Semanal 2002;Semana epidemiológica No. 2.

26. SIVIGILA. Enfermedades de notificación obligatoria. Informe primer semestre de 2003. Boletín Epidemiológico Semanal 2003;Semana epidemiológica No. 26.
27. Ocazonez RE, Gomez SY, Cortes FM. [Dengue hemorrhagic fever serotype and infection pattern in a Colombian endemic area]. *Revista de salud publica* 2007;**9**:262–74.
28. WHO. Dengue haemorrhagic fever: Diagnosis, treatment, prevention and control. World Health Organization; 1997.
29. Conroy AL, Gelvez M, Hawkes M, et al. Host biomarkers distinguish dengue from leptospirosis in Colombia: a case-control study. *BMC infectious diseases* 2014;**14**:35.
30. Khongphatthanayothin A, Phumaphuti P, Thongchaiprasit K, Poovorawan Y. Serum levels of sICAM-1 and sE-selectin in patients with dengue virus infection. *Jpn J Infect Dis* 2006;**59**:186–8.
31. Cardier JE, Rivas B, Romano E, et al. Evidence of vascular damage in dengue disease: demonstration of high levels of soluble cell adhesion molecules and circulating endothelial cells. *Endothelium* 2006;**13**:335–40.
32. Levy A, Valero N, Espina LM, Anez G, Arias J, Mosquera J. Increment of interleukin 6, tumour necrosis factor alpha, nitric oxide, C-reactive protein and apoptosis in dengue. *Trans R Soc Trop Med Hyg* 2010;**104**:16–23.
33. de-Oliveira-Pinto LM, Gandini M, Freitas LP, et al. Profile of circulating levels of IL-1Ra, CXCL10/IP-10, CCL4/MIP-1beta and CCL2/MCP-1 in dengue fever and parvovirus. *Mem Inst Oswaldo Cruz* 2012;**107**:48–56.
34. Becerra A, Warke RV, Martin K, et al. Gene expression profiling of dengue infected human primary cells identifies secreted mediators in vivo. *J Med Virol* 2009;**81**:1403–11.
35. Fink J, Gu F, Ling L, et al. Host gene expression profiling of dengue virus infection in cell lines and patients. *PLoS Negl Trop Dis* 2007;**1**:e86.
36. Michels M, van der Ven AJ, Djamiatun K, et al. Imbalance of Angiopoietin-1 and Angiopoietin-2 in Severe Dengue and Relationship with Thrombocytopenia, Endothelial Activation, and Vascular Stability. *Am J Trop Med Hyg* 2012.
37. Brasier AR, Ju H, Garcia J, et al. A three-component biomarker panel for prediction of dengue hemorrhagic fever. *Am J Trop Med Hyg* 2012;**86**:341–8.
38. Ju H, Brasier AR. Variable selection methods for developing a biomarker panel for prediction of dengue hemorrhagic fever. *BMC research notes* 2013;**6**:365.
39. Malavice GN, Gomes L, Alles L, et al. Serum IL-10 as a marker of severe dengue infection. *BMC infectious diseases* 2013;**13**:341.
40. Srikiathachorn A, Ajariyakhajorn C, Endy TP, et al. Virus-induced decline in soluble vascular endothelial growth receptor 2 is associated with plasma leakage in dengue hemorrhagic fever. *J Virol* 2007;**81**:1592–600.
41. Low JG, Ong A, Tan LK, et al. The early clinical features of dengue in adults: challenges for early clinical diagnosis. *PLoS Negl Trop Dis* 2011;**5**:e1191.
42. Lee CG, Da Silva CA, Dela Cruz CS, et al. Role of chitin and chitinase/chitinase-like proteins in inflammation, tissue remodeling, and injury. *Annu Rev Physiol* 2011;**73**:479–501.
43. Nordenbaek C, Johansen JS, Junker P, Borregaard N, Sorensen O, Price PA. YKL-40, a matrix protein of specific granules in neutrophils, is elevated in serum of patients with community-acquired pneumonia requiring hospitalization. *J Infect Dis* 1999;**180**:1722–6.
44. Kronborg G, Ostergaard C, Weis N, et al. Serum level of YKL-40 is elevated in patients with Streptococcus pneumoniae bacteremia and is associated with the outcome of the disease. *Scandinavian journal of infectious diseases* 2002;**34**:323–6.
45. Erdman LK, Petes C, Lu Z, et al. Chitinase 3-like 1 is induced by Plasmodium falciparum malaria and predicts outcome of cerebral malaria and severe malarial anaemia in a case-control study of African children. *Malar J* 2014;**13**:279.
46. Fagundes CT, Costa VV, Cisalpino D, et al. IFN-gamma production depends on IL-12 and IL-18 combined action and mediates host resistance to dengue virus infection in a nitric oxide-dependent manner. *PLoS Negl Trop Dis* 2011;**5**:e1449.
47. Camenisch G, Pisabarro MT, Sherman D, et al. ANGPTL3 stimulates endothelial cell adhesion and migration via integrin alpha vbeta 3 and induces blood vessel formation in vivo. *The Journal of biological chemistry* 2002;**277**:17281–90.
48. Ono M, Shimizugawa T, Shimamura M, et al. Protein region important for regulation of lipid metabolism in angiopoietin-like 3 (ANGPTL3): ANGPTL3 is cleaved and activated in vivo. *The Journal of biological chemistry* 2003;**278**:41804–9.
49. Shimamura M, Matsuda M, Kobayashi S, et al. Angiopoietin-like protein 3, a hepatic secretory factor, activates lipolysis in adipocytes. *Biochemical and biophysical research communications* 2003;**301**:604–9.
50. Botella LM, Sanchez-Elsner T, Sanz-Rodriguez F, et al. Transcriptional activation of endoglin and transforming growth factor-beta signaling components by cooperative interaction between Sp1 and KLF6: their potential role in the response to vascular injury. *Blood* 2002;**100**:4001–10.
51. Torsney E, Charlton R, Parums D, Collis M, Arthur HM. Inducible expression of human endoglin during inflammation and wound healing in vivo. *Inflammation research: official journal of the European Histamine Research Society [et al]* 2002;**51**:464–70.
52. Hawinkels LJ, Kuiper P, Wiercinska E, et al. Matrix metalloproteinase-14 (MT1-MMP)-mediated endoglin shedding inhibits tumor angiogenesis. *Cancer research* 2010;**70**:4141–50.
53. Venkatesha S, Toporsian M, Lam C, et al. Soluble endoglin contributes to the pathogenesis of preeclampsia. *Nat Med* 2006;**12**:642–9.
54. Dietmann A, Helbok R, Lackner P, et al. Endoglin in African children with Plasmodium falciparum malaria: a novel player in severe malaria pathogenesis? *J Infect Dis* 2009;**200**:1842–8.
55. Erdman LK, Dhabangi A, Musoke C, et al. Combinations of Host Biomarkers Predict Mortality among Ugandan Children with Severe Malaria: A Retrospective Case-Control Study. *PLoS One* 2011;**6**:e17440.
56. Silver KL, Conroy AL, Leke RG, et al. Circulating soluble endoglin levels in pregnant women in cameroon and Malawi-associations with placental malaria and fetal growth restriction. *PLoS One* 2011;**6**:e24985.
57. Levine RJ, Lam C, Qian C, et al. Soluble endoglin and other circulating anti-angiogenic factors in preeclampsia. *N Engl J Med* 2006;**355**:992–1005.
58. van de Weg CA, Pannuti CS, van den Ham HJ, et al. Serum angiopoietin-2 and soluble VEGF receptor 2 are surrogate markers for plasma leakage in patients with acute dengue virus infection. *J Clin Virol* 2014;**60**:328–35.
59. Brown MG, McAlpine SM, Huang YY, et al. RNA sensors enable human mast cell anti-viral chemokine production and IFN-mediated protection in response to antibody-enhanced dengue virus infection. *PLoS One* 2012;**7**:e34055.
60. Warke RV, Becerra A, Zawadzka A, et al. Efficient dengue virus (DENV) infection of human muscle satellite cells upregulates type I interferon response genes and differentially modulates MHC I expression on bystander and DENV-infected cells. *The Journal of general virology* 2008;**89**:1605–15.
61. Tolfvenstam T, Lindblom A, Schreiber MJ, et al. Characterization of early host responses in adults with dengue disease. *BMC infectious diseases* 2011;**11**:209.
62. Chen JP, Lu HL, Lai SL, et al. Dengue virus induces expression of CXC chemokine ligand 10/IFN-gamma-inducible protein 10, which competitively inhibits viral binding to cell surface heparan sulfate. *J Immunol* 2006;**177**:3185–92.
63. Ip PP, Liao F. Resistance to dengue virus infection in mice is potentiated by CXCL10 and is independent of CXCL10-mediated leukocyte recruitment. *J Immunol* 2010;**184**:5705–14.
64. Kelley JF, Kaufusi PH, Volper EM, Nerurkar VR. Maturation of dengue virus nonstructural protein 4B in monocytes enhances production of dengue hemorrhagic fever-associated chemokines and cytokines. *Virology* 2011;**418**:27–39.
65. Hsieh MF, Lai SL, Chen JP, et al. Both CXCR3 and CXCL10/IFN-inducible protein 10 are required for resistance to primary infection by dengue virus. *J Immunol* 2006;**177**:1855–63.
66. Becquart P, Wauquier N, Nkoghe D, et al. Acute dengue virus 2 infection in Gabonese patients is associated with an early innate immune response, including strong interferon alpha production. *BMC infectious diseases* 2010;**10**:356.
67. Salgado DM, Eltit JM, Mansfield K, et al. Heart and skeletal muscle are targets of dengue virus infection. *Pediatr Infect Dis J* 2010;**29**:238–42.
68. Wang WK, Chen HL, Yang CF, et al. Slower rates of clearance of viral load and virus-containing immune complexes in patients with dengue hemorrhagic fever. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America* 2006;**43**:1023–30.
69. Libraty DH, Endy TP, Hough HS, et al. Differing influences of virus burden and immune activation on disease severity in secondary dengue-3 virus infections. *J Infect Dis* 2002;**185**:1213–21.
70. Vaughn DW, Green S, Kalayanaraj S, et al. Dengue viremia titer, antibody response pattern, and virus serotype correlate with disease severity. *J Infect Dis* 2000;**181**:2–9.
71. Cardier JE, Marino E, Romano E, et al. Proinflammatory factors present in sera from patients with acute dengue infection induce activation and apoptosis of human microvascular endothelial cells: possible role of TNF-alpha in endothelial cell damage in dengue. *Cytokine* 2005;**30**:359–65.
72. Lin CF, Chiu SC, Hsiao YL, et al. Expression of cytokine, chemokine, and adhesion molecules during endothelial cell activation induced by antibodies against dengue virus nonstructural protein 1. *J Immunol* 2005;**174**:395–403.
73. Brown MG, Hermann LL, Issekutz AC, et al. Dengue virus infection of mast cells triggers endothelial cell activation. *J Virol* 2011;**85**:1145–50.
74. Kelley JF, Kaufusi PH, Nerurkar VR. Dengue hemorrhagic fever-associated immunomodulators induced via maturation of dengue virus nonstructural 4B protein in monocytes modulate endothelial cell adhesion molecules and human microvascular endothelial cells permeability. *Virology* 2012;**422**:326–37.
75. Azeredo EL, Zagne SM, Alvarenga AR, Nogueira RM, Kubelka CF, de Oliveira-Pinto LM. Activated peripheral lymphocytes with increased expression of cell adhesion molecules and cytotoxic markers are associated with dengue fever disease. *Mem Inst Oswaldo Cruz* 2006;**101**:437–49.
76. Azeredo EL, Neves-Souza PC, Alvarenga AR, et al. Differential regulation of toll-like receptor-2, toll-like receptor-4, CD16 and human leucocyte antigen-DR on peripheral blood monocytes during mild and severe dengue fever. *Immunology* 2010;**130**:202–16.
77. Valero N, Larreal Y, Espina LM, Reyes I, Maldonado M, Mosquera J. Elevated levels of interleukin-2 receptor and intercellular adhesion molecule 1 in sera from a venezuelan cohort of patients with dengue. *Arch Virol* 2008;**153**:199–203.
78. Bethell DB, Flobbe K, Cao XT, et al. Pathophysiologic and prognostic role of cytokines in dengue hemorrhagic fever. *J Infect Dis* 1998;**177**:778–82.
79. Villar-Centeno LA, Diaz-Quijano FA, Martinez-Vega RA. Biochemical alterations as markers of dengue hemorrhagic fever. *Am J Trop Med Hyg* 2008;**78**:370–4.
80. Diaz-Quijano FA, Martinez-Vega RA, Villar-Centeno LA. Early predictors of haemorrhage in acute febrile syndrome patients from Bucaramanga, Colombia: a dengue endemic area. *Singapore medical journal* 2008;**49**:480–2.
81. Shepard DS, Undurraga EA, Betancourt-Cravioto M, et al. Approaches to refining estimates of global burden and economics of dengue. *PLoS Negl Trop Dis* 2014;**8**:e3306.
82. Suaya JA, Shepard DS, Siqueira JB, et al. Cost of dengue cases in eight countries in the Americas and Asia: a prospective study. *Am J Trop Med Hyg* 2009;**80**:846–55.

83. Guilarde AO, Turchi MD, Siqueira Jr JB, et al. Dengue and dengue hemorrhagic fever among adults: clinical outcomes related to viremia, serotypes, and antibody response. *J Infect Dis* 2008;**197**:817–24.
84. Ozturk B, Kuscu F, Tutuncu E, Sencan I, Gurbuz Y, Tuzun H. Evaluation of the association of serum levels of hyaluronic acid, sICAM-1, sVCAM-1, and VEGF-A with mortality and prognosis in patients with Crimean-Congo hemorrhagic fever. *J Clin Virol* 2010;**47**:115–9.
85. Zhang Y, Liu B, Ma Y, et al. Hantaan virus infection induces CXCL10 expression through TLR3, RIG-I, and MDA-5 pathways correlated with the disease severity. *Mediators of inflammation* 2014;**2014**:697837.
86. Wang PZ, Li ZD, Yu HT, et al. Elevated serum concentrations of inflammatory cytokines and chemokines in patients with haemorrhagic fever with renal syndrome. *The Journal of international medical research* 2012;**40**:648–56.
87. Connolly-Andersen AM, Thunberg T, Ahlm C. Endothelial activation and repair during hantavirus infection: association with disease outcome. *Open forum infectious diseases* 2014;**1**:ofu027.
88. ter Meulen J, Sakho M, Koulemou K, et al. Activation of the cytokine network and unfavorable outcome in patients with yellow fever. *J Infect Dis* 2004;**190**:1821–7.
89. Wauquier N, Becquart P, Padilla C, Baize S, Leroy EM. Human fatal zaire ebola virus infection is associated with an aberrant innate immunity and with massive lymphocyte apoptosis. *PLoS Negl Trop Dis* 2010;**4**.
90. McElroy AK, Erickson BR, Flietstra TD, et al. Biomarker correlates of survival in pediatric patients with Ebola virus disease. *Emerging infectious diseases* 2014;**20**:1683–90.
91. McElroy AK, Erickson BR, Flietstra TD, et al. Ebola hemorrhagic Fever: novel biomarker correlates of clinical outcome. *J Infect Dis* 2014;**210**:558–66.